

Referências bibliográficas LÂMINA LANÇAMENTO ESPESSADO x RR

- 1.Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
- 2.Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2011;52:238–225
- 3.Boehm G et al. Prebiotics in infant formulas. *J Clin Gastroenterol.* 2004; 38: S76-9.
- 4.Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. *Br J Nutr.* 2015;113(9):1339-44.
- 5.Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
6. Comparativo de tabela nutricional de produtos da mesma categoria realizado em Agosto 2024.
- 7.Salvatore, S. et al. Thickened infant formula: what to know. *Nutrition*, 2018.
- 8.Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2018;66(3):516-554.
- 9.Infante Pina D et al. Thickened infant formula, rheological study of the “in vitro” properties. *An Pediatr (Barc).* 2010; 72(5):302-8.
- 10.Horvath A, Dziechciarz P, Szajewska H. The Effect of Thickened-Feed Interventions on Gastroesophageal Reflux in Infants: Systematic Review and Meta-analysis of Randomized, Controlled Trials. *Pediatrics* 2008;122:e1268–e1277.
- 11.Koletzko B, Karin B, Thomas BJ, Phipip C Should formula for infants provide arachidonic acid along with DHA? A position paper of the European Academy of Paediatrics and the Child Health Foundation. *Am J Clin Nutri.* 2020. Vol. 111, nº 1, p. 10-16.
- 12.Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
- 13.Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics.* 2013;131(1):262-72.
- 14.Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
- 15.Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients.* 2021; 13: 247.

16. Lerner A, Shamir R. Nucleotides in infant nutrition: a must or an option. *Isr Med Assoc J.* 2000; 2(10):772-4.
17. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
18. Dupont C. Protein requirements during the first year of life. *Am J Clin Nutr.* 2003;77(6):1544S-1549S.
19. EFSA. Scientific Opinion on the essential composition of infant and follow-on formulae. *EFSA Journal.* 2014;12(7).
20. Sociedade Brasileira de Pediatria. Departamento de Nutrologia. Manual de Alimentação: Da Infância à adolescência, 2018.
21. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol,* 1997; 29:237-42.
22. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics,* 2003;111(4):355-359
23. Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. *Pediatr. Mod.* 2014; 50(8): 339-342.
24. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. *Acta Pædiatrica* 2015, ISSN 0803-5253
25. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. *Eur J Clin Nutr* 2006, 60:1304-10.
26. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr Suppl* 2003, 91:86-90.
27. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr* 2003, 36:343-51.
28. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. *Nutr J* 2007;6:8.
29. Savino F et al. Advances in the management of digestive problems during the first months of life. *Acta Paediatr Suppl* 2005, 94:120.
30. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai.* 2004 Jun;87(6):641- 9.