

REFERÊNCIAS BIBLIOGRÁFICAS

Material Aptamil Pepti HMO Lançamento

- 1) HCP TRACKER, 2024
- 2) Epifanio, M. et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. *Biomed J Sci & Tech Res* 25(1)-2020. BJSTR. MS.ID.00416.
- 3) Vandenplas Y, et al. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. *J Pediatr Gastroenterol Nutr.* 2024 Feb;78(2):386-413.
- 4) *Nutrologia Pediátrica: Temas da Atualidade em Nutrologia Pediátrica - 2021.* / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2021.
- 5) Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.*2013;60(1):49-74
- 6) FERREIRA, CHT; BARRETO, BAP; TOFOLI, MHC; et al. *Microbioma e pediatria na saúde e na doença.* Barueri [SP]: Manole, 2024. 1.ed.
- 7) *Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar/ Sociedade Brasileira de Pediatria. – 5.ed. rev. ampl.– São Paulo: SBP, 2024. 208 p.*
- 8) Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *J Biol Regul Homeost Agents.* 2012;26(3 Suppl):49-59.
- 9) Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr.* 2008;138(6):1091-5.
- 10) Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child.* 2006;91(10):814-
- 11) Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
- 12) Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr.* 2005;41(5):584-99

- 13) SALMINEN, Seppo; SZAJEWSKA, Hania; KNOL, Jan. *The Biotics Family in Early Life*. 2. ed. Wiley, 2023.
- 14) Goh CY, et al. Combination of short-chain GOS and long-chain FOS 9:1 with 2FL positively impact the infant gut microbiota composition and metabolic activity in a stimulator of the human intestinal microbial ecosystem (SHIME). *J Pediatr Gastroenterol Nutr* 2019; 68 (S1): NP 114; 1159.
- 15) Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol*. 2018;2(1):7-82.
- 16) Vieira MC, Morais MB, Spolidoro JVN, et al. A survey on clinical presentation and nutritional status of infants with suspected cow's milk allergy. *BMC Pediatr*. 2010;10:1-7.
- 17) World Allergy Organization (WAO) Diagnosis and Rationale for Action against Cow's Milk Allergy (DRACMA) guidelines update – XVI - Nutritional management of cow's milk allergy Venter, Carina Ansotegui, Ignacio et al. *World Allergy Organization Journal*, Volume 17, Issue 8, 2024.
- 18) Vandenplas Y, et al. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. *J Pediatr Gastroenterol Nutr*. 2024 Feb;78(2):386-413.
- 19) Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
- 20) Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr*. 1995;49 Suppl 1:S39 S48.2.
- 21) Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol*. 2001;12(2):83 86.
- 22) Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics*, 90(3), 463 465.4.
- 23) Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics*, 127(4), 550 557.
- 24) SBP. Guia de orientações - Dificuldades alimentares/ Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia . São Paulo: SBP, 2022. 66 f.
- 25) Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol*. 2012;23(4):307-14.